

2025



VETERINARY
DIAGNOSTICS
Catalogue



TABLE OF CONTENTS

FARM LIVESTOCK	2
Cow and Goat Pregnancy Tests	2
Livestock Infections	3
PETS	5
Infections	5
Hormones	7
Allergy	8

Kit format: 96-well microplate, breakable strips, all reagents are liquid and ready-to-use, dispensed in plastic vials with screw caps, chromogen – single component TMB, shelf life – 18 months.

XEMA

Xema Oy, Finland

✉ info@xema.fi

🌐 www.xema.fi

FARM LIVESTOCK

Cow and Goat Pregnancy Tests

[REF] K207M Progesterone milk EIA (Cows pregnancy test)

A solid-phase enzyme immunoassay for the quantitative determination of progesterone in cow's milk

Sample type:	cow's milk	Incubation:	60'/15'; 37 °C	Control sample:	1
Sample volume:	25 µl	Calibrators:	6 (0-160 ng/ml)	Shelf life:	18 months
Sensitivity:	0.3 ng/ml				

[REF] K247M Milk PAG EIA

A solid-phase enzyme immunoassay for a semi-quantitative determination of Pregnancy-Associated Globulin (PAG) in cow's milk

Sample type:	cow's milk	Incubation:	120'/15'; 37°C, shaker	Control sample:	2
Sample volume:	100 µl	Calibrators:	2	Shelf life:	18 months

[REF] K247B PAG EIA

A solid-phase enzyme immunoassay for a semi-quantitative determination of Pregnancy-Associated Glycoproteins (PAG) in cow and goat serum

Sample type:	serum	Incubation:	30'/30'/15'; 37°C, shaker	Control sample:	2
Sample volume:	50 µl	Calibrators:	2	Shelf life:	18 months

[REF] K271B Colostrum IgG EIA

New test!

A solid-phase enzyme immunoassay for the quantitative determination of IgG in bovine colostrum

Sample type:	colostrum	Sample volume:	50 µl	Calibrators:	5 (0-100 µg/ml)
Sample predilution:	1:401	Incubation:	60'/15'; 37°C	Shelf life:	18 months

Livestock Infections

[REF] K107B Anti-Bovine Leukemia Virus IgG EIA

A solid-phase enzyme immunoassay for the qualitative determination of BLV antigen in bovine serum or milk

Sample type:	serum, plasma	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:41	Incubation:	60'/60'/15'; 37°C	Shelf life:	18 months

[REF] K116B Brucella IgG-B EIA

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to *Brucella* in bovine serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:41	Incubation:	60'/60'/15'; 37°C	Shelf life:	18 months

[REF] K157S Ab to Swine Fever Virus EIA

New test!

A solid-phase enzyme immunoassay for the qualitative determination of antibodies to Swine Fever Virus in serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Calibrators:	2
Sample predilution:	1:100	Incubation:	30'/30'/15'; 37°C	Shelf life:	18 months

[REF] K106Y MG Ab EIA

A solid-phase enzyme immunoassay for the qualitative determination of antibodies to *Mycoplasma gallisepticum* in chicken and turkey serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:501	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

[REF] K150Y Influenza A Ab EIA

A solid-phase enzyme immunoassay for the qualitative determination of antibodies to Influenza A virus in chicken and turkey serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:501	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

[REF] K152Y NDV IgG EIA

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to Newcastle Disease virus (NDV) in chicken or turkey serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Control samples:	2
Sample dilution:	1:501	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

[REF] K153Y IBV Ab EIA

A solid-phase enzyme immunoassay for the qualitative determination of antibodies to Avian Infectious Bronchitis virus (IBV) in birds' serum

Sample type:	serum	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:501	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

[REF] K154Y EDS-76 Virus Ab EIA

New test!

A solid-phase enzyme immunoassay for the qualitative determination of antibodies to Egg Drop Syndrome-76 (EDS-76) virus in chicken and turkey serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:501	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

REF K155Y APV Ab EIA

A solid-phase enzyme immunoassay for the qualitative determination of antibodies to Avian Pneumovirus (APV) in birds' serum

Sample type:	serum	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:501	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

REF K165Y Reo Ab EIA

A solid-phase enzyme immunoassay for the qualitative determination of antibodies to Avian Reovirus (ARV) in birds' serum

Sample type:	serum	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:501	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

REF K166Y IBD Ab EIA

A solid-phase enzyme immunoassay for the qualitative determination of antibodies to Infectious Bursal Disease virus (IBDV) in birds' serum

Sample type:	serum	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:501	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

REF K184Y ABP Ab EIA***New test!***

A solid-phase enzyme immunoassay for the qualitative determination of antibodies to Avibacterium (Haemophilus) paragallinarum (ABP) in chicken or turkey serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:501	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

REF K188Y SE Ab EIA***New test!***

A solid-phase enzyme immunoassay for the qualitative determination of antibodies to *Salmonella enteritidis* in chicken and turkey serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:501	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

REF K189Y PM Ab EIA***New test!***

A solid-phase enzyme immunoassay for the qualitative determination of antibodies to *Pasteurella multocida* in chicken and turkey serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:501	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

PETS**Infections****REF K101CF Toxoplasma IgG-CF EIA**

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to *Toxoplasma gondii* in canine or feline serum or plasma

Sample type:	serum, plasma	Incubation:	30'/30'/15'; 37 °C	Control samples:	2
Sample volume:	100 µl	Calibrator:	1	Shelf life:	18 months
Sample predilution:	1:21				

REF K104CF HSV 1/2 IgG-CF EIA

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to HSV 1/2 in canine or feline serum or plasma

Sample type:	serum, plasma	Incubation:	30'/30'/15'; 37 °C	Control samples:	2
Sample volume:	100 µl	Calibrator:	1	Shelf life:	18 months
Sample predilution:	1:21				

REF K105CF Chlamydia IgG-CF EIA

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to *Chlamydia* in canine or feline serum or plasma

Sample type:	serum, plasma	Incubation:	30'/30'/15'; 37 °C	Control samples:	2
Sample volume:	100 µl	Calibrator:	1	Shelf life:	18 months
Sample predilution:	1:21				

REF K106CF Mycoplasma IgG-CF EIA

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to *Mycoplasma* in canine or feline serum or plasma

Sample type:	serum, plasma	Incubation:	30'/30'/15'; 37 °C	Control samples:	2
Sample volume:	100 µl			Shelf life:	18 months
Sample predilution:	1:21				

REF K112F Feline Immunodeficiency Virus IgG EIA

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to feline immunodeficiency virus in feline serum or plasma

Sample type:	serum, plasma	Incubation:	60'/60'/15'; 37 °C	Shelf life:	18 months
Sample volume:	10 µl	Control samples:	2		

REF K116C Brucella IgG-C EIA

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to *Brucella* in canine serum or plasma

Sample type:	serum, plasma	Incubation:	30'/15'/15'; 18-25°C,	Shelf life:	18 months
Sample volume:	20 µl	Controls:	shaker 2		

REF K122C Candida IgG-C EIA

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to *Candida* in canine serum or plasma

Sample type:	serum, plasma	Incubation:	30'/30'/15'; 37 °C	Control samples:	2
Sample volume:	100 µl	Calibrator:	1	Shelf life:	18 months
Sample predilution:	1:101				

REF K151C Parvo-IgG-C EIA

A solid-phase enzyme immunoassay for the quantitative determination of Parvo-IgG in serum or plasma of *Canidae* and *Mustelidae*

Sample type:	serum, plasma	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months
Sample volume:	100 µl	Calibrators:	6 (0–320 U/ml)		
Sample predilution:	1:101				

REF K152C CDV-IgG-C EIA

A solid-phase enzyme immunoassay for the quantitative determination of IgG antibodies to CDV in serum or plasma of *Canidae* and *Mustelidae*

Sample type:	serum, plasma	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months
Sample volume:	100 µl	Calibrators:	6 (0–320 U/ml)		
Sample predilution:	1:101				

REF K1532F Feline SARS-Cov-2 antibody EIA

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to SARS-Cov-2 in feline serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Control samples:	3
Sample dilution:	1:101	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

REF X153F XEMAtest Corona F

Rapid immunochromatographic test for the qualitative determination of antibodies to SARS-CoV-2 in feline whole blood, serum, plasma or ascitic fluid

Sample type:	whole blood, serum, plasma, ascitic fluid	Sample volume:	40 µl	Shelf life:	24 months
		Incubation:	10', 18–25 °C		

REF K159CF Rabies Virus IgG-CF EIA

A solid-phase enzyme immunoassay for the quantitative determination of IgG antibodies to Rabies virus in canine and feline serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Controls:	2
Sample predilution:	1:50	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

REF K159P Rabies Virus IgG-P EIA**New test!**

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to Rabies Virus in Primates serum or plasma

Sample type:	serum, plasma	Sample volume:	100 µl	Controls:	2
Sample predilution:	1:50	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

REF K059 Rabies Virus EIA

A solid-phase enzyme immunoassay for the quantitative determination of Rabies Virus Antigen in culture fluid

Sample type:	Culture fluid	Sample volume:	100 µl	Control sample:	no
Calibrators:	7 (0 – 5 IU/ml)	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months
Sensitivity:	0,07 IU/ml				

REF K174CF Nematoda IgG-CF EIA

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to *Nematoda* in serum or plasma of *Carnivora*

Sample type:	serum, plasma	Sample predilution:	1:101	Control samples:	2
Sample volume:	100 µl	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

REF K175CF Eucestoda IgG-CF EIA

A solid-phase enzyme immunoassay for the qualitative determination of IgG antibodies to *Eucestoda* in serum or plasma of *Carnivora*

Sample type:	serum, plasma	Sample volume:	100 µl	Control samples:	2
Sample predilution:	1:101	Incubation:	30'/30'/15'; 37 °C	Shelf life:	18 months

REF K250C Canine CRP EIA

Sample type:	serum, plasma	Incubation:	30'/30'/15'; 37 °C	Control samples:	1
Sample predilution:	1:5000	Calibrators:	5(0-500) mg/l	Shelf life:	18 months
Sample volume:	100 µl				

Hormones

REF K201C Canine TSH EIA

A solid-phase enzyme immunoassay for the quantitative determination of canine TSH in canine serum (plasma)

Sample type:	serum, plasma	Incubation:	120'/15'; 37 °C	Control sample:	1
Sample volume:	50 µl	Calibrators:	5 (0–8 ng/ml)	Shelf life:	18 months
Sensitivity:	0.05 ng/ml				

REF K211CF total T3-CF EIA

A solid-phase enzyme immunoassay for the quantitative determination of total T3 in canine or feline serum or plasma

Sample type:	serum or plasma	Incubation:	60'/15'; 37 °C	Control sample:	1
Sample volume:	25 µl	Calibrators:	5 (0–15 nmol/l)	Shelf life:	18 months
Sensitivity:	0.4 nmol/l				

REF K212CF total T4-CF EIA

A solid-phase enzyme immunoassay for the quantitative determination of total T4 in canine or feline serum or plasma

Sample type:	serum or plasma	Incubation:	60'/15'; 37 °C	Control sample:	1
Sample volume:	25 µl	Calibrators:	5 (0–160 nmol/l)	Shelf life:	18 months
Sensitivity:	6.0 nmol/l				

REF K213CF free T3-CF EIA

A solid-phase enzyme immunoassay for the quantitative determination of free T3 in canine or feline serum or plasma

Sample type:	serum or plasma	Incubation:	60'/15'; 37 °C	Control sample:	1
Sample volume:	25 µl	Calibrators:	6 (0–40 pmol/l)	Shelf life:	18 months
Sensitivity:	1.0 pmol/l				

REF K214CF free T4-CF EIA

A solid-phase enzyme immunoassay for the quantitative determination of free T4 in canine or feline serum or plasma

Sample type:	serum or plasma	Incubation:	60'/15'; 37 °C	Control sample:	1
Sample volume:	25 µl	Calibrators:	6 (0–100 pmol/l)	Shelf life:	18 months
Sensitivity:	1.0 pmol/l				

REF K202C Canine LH EIA

Sample type:	serum, plasma	Incubation:	60'/15'; 37 °C	Control sample:	no
Sample volume:	50 µl	Calibrators:	4 (0–50 ng/ml)	Shelf life:	18 months

REF K207CF Progesterone-CF EIA

A solid-phase enzyme immunoassay for the quantitative determination of Progesterone in canine or feline serum or plasma

Sample type:	serum, plasma	Incubation:	120'/15'; 37 °C	Control sample:	1
Sample volume:	25 µl	Calibrators:	7 (0–300 nmol/l)	Shelf life:	18 months
Sensitivity:	0.5 nmol/l				

REF K208CF Estradiol-CF EIA

Sample type:	serum, plasma	Incubation:	120'/15'; 37°C	Control sample:	1
Sample volume:	25 µl	Calibrators:	5 (0-300 pg/ml)	Shelf life:	18 months
Sensitivity:	1.0 pg/ml				

REF K209CF Testosterone-CF EIA

Sample type:	serum, plasma	Incubation:	120'/15'; 37°C	Control sample:	1
Sample volume:	25 µl	Calibrators:	7 (0-100 nmol/l)	Shelf life:	18 months
Sensitivity:	0.25 nmol/l				

REF K210CF Cortisol-CF EIA

A solid-phase enzyme immunoassay for the quantitative determination of Cortisol in canine or feline serum or plasma

Sample type:	serum, plasma	Incubation:	60'/15'; 37 °C	Control sample:	1
Sample volume:	10 µl	Calibrators:	6 (0–600 nmol/l)	Shelf life:	18 months
Sensitivity:	10 nmol/l				

REF K210R Rat/Mouse Corticosterone EIA

Sample type:	serum, plasma	Incubation:	60'/15'; 37°C	Control sample:	1
Sample volume:	25 µl	Calibrators:	6(0-1000) nmol/l	Shelf life:	18 months
Sensitivity:	5.0 nmol/l				

Allergy

REF K200C Canine IgE EIA

Sample type:	serum, plasma	Sample volume:	10 µl	Calibrators:	6 (0-280) mg/l
Sample predilution:	1:101	Incubation:	60'/15'; 37°C	Shelf life:	18 months

Xema Oy (Finland)

Myllymaenkatu, 21, 53550, Lappeenranta;

Tel./fax: +358 401 922 737

info@xema.fi

www.xema.fi



CE

